arXiv:math/0507232v3 [math.AG] 25 Jul 2007

Quantum cohomology of smooth complete intersections in weighted projective spaces and singular toric varieties

VICTOR PRZYJALKOWSKI

Abstract. We generalize Givental's Theorem for complete intersections in smooth toric varieties in the Fano case. In particular, we find Gromov–Witten invariants of Fano varieties of dimension ≥ 3, which are complete intersections in weighted projective spaces and singular toric varieties. We generalize the Riemann–Roch equations to such varieties. We compute counting matrices of smooth Fano threefolds with Picard group $\mathbb{Z}$ and anticanonical degrees 2, 8, and 16.

Let X be a general complete intersection of hypersurfaces of degrees $d_1, \ldots, d_l$ in $\mathbb{P}^N$, $d_0 = N+1-\sum d_i > 0$, $n = N - l \geqslant 3$. Let $H^*_H(X) \subset H^*(X, \mathbb{Z})$ be the ring generated by class H that is dual to the hyperplane section and $S = H^*_H(X) \otimes \mathbb{C}[[q]]$. Denote $H \otimes 1 \in S$, $H \in H^*_H(X)$ by the same letter H. According to Givental's Theorem for complete intersections in projective spaces (see [Gi96]), I-series for X (i.e. the generating series for one-pointed Gromov–Witten invariants with descendants, see definition 2.2.3) is

$$I^X = e^{-\alpha_X q} \sum_{d=0}^{\infty} \widehat{I}^X_d q^d = e^{-\alpha_X q} \sum_{d=0}^{\infty} q^d \frac{\prod_{i=1}^{l} \prod_{j=1}^{dd_i} (d_i H + j)}{\prod_{i=1}^{d} (H+i)^{N+1}} \in S,$$

where $\alpha_X = d_1! \ldots d_l!$ if $d_0 = 1$ and $\alpha_X = 0$ if $d_0 > 1$. Consider $D = q \cdot d/dq \in \mathcal{D} = \mathbb{C}[[q]][q \cdot d/dq]$. Then a solution of the Riemann–Roch equation (or equation of type Dn), i. e. the differential equation

$$(D^{N+1} - \prod_{i=0}^{l} d_i^{d_i} q \prod_{j=1}^{d_i} (D + \frac{j}{d_i}))[\Phi(q)] = 0$$

is the series $\mathcal{I}^X = \sum q^d (d_0 d)! \widehat{I}^X_{d,H^0}$.

In this paper we generalize these results to complete intersections in *weighted projective spaces* and *singular toric varieties.*

In the first section we formulate the main results. In the second section we give definitions and formulate known results. In the third section we prove and discuss the main theorems 1.1 and 1.2. In Appendix we formulate Golyshev's conjecture. The corollary of theorem 1.1 completes its proof.

Notations. *The Pochhammer symbol* $\boldsymbol{(X)_n}$ denotes the product $X(X+1) \cdot \ldots \cdot (X+n-1)$, where X belongs to some ring (see [AS72], 6.1.22). An infinite product of the form $\prod_{a=-\infty}^{n-1} (X+a)$ is denoted by $\boldsymbol{[X]_n}$ (we define the infinite product formally; we will only use quotients of two such products in which all but for a finite number of factors coincide).

We denote the groups of homological and cohomological classes on X modulo torsion by $H_*(X)$ and $H^*(X)$ accordingly.

The work was partially supported by RFFI grants $04-01-00613$ and $05-01-00353$, and grant NSh$-$9969.2006.1.

We use the same notation for a hypersurface and its dual cohomological class.
Denote $B = \mathbb{C}[[q]]$ and $D = q \cdot d/dq \in \mathcal{D} = B[q \cdot d/dq]$.

Everything is over $\mathbb{C}$.

1. RESULTS

See 2.1, 2.2 and 2.4 for necessary definitions.

1.1. **Theorem.** *Let $\mathbb{P} = \mathbb{P}(w_1, \ldots, w_k)$ be a weighted projective space. Let X be a smooth complete intersection of hypersurfaces $X_1, \ldots, X_l$ which do not intersect the singular locus of $\mathbb{P}$. Assume that $-K_X > 0$ and $\operatorname{Pic} X = \mathbb{Z}H$, where H is the class dual to the hyperplane. Let $i\colon X \to \mathbb{P}$ be the natural embedding.*

1) *I-series for X is the following.*

$$I^X = e^{-\alpha_X q} \sum_{d=0}^{\infty} q^d \cdot i^* \left(\frac{\prod_{a=1}^{l} \big(X_a + 1\big)_{\boldsymbol{d \cdot \deg X_a}}}{\prod_{a=1}^{k} \big(w_a H + 1\big)_{\boldsymbol{d \cdot w_a}}} \right).$$

Here $\alpha_X = 0$ if the index of X is 2 or greater and $\alpha_X = \prod_{a=1}^{l} (\deg X_a)!/\prod_{a=1}^{k} w_a!$ if the index is 1.

2) *Let d_i $(1 \le i \le l)$ be the degrees (with respect to H) of the hypersurfaces X_i and $d_0 = \sum w_i - \sum d_i$ be the index of X.Let $\mathcal{I}^X(q) = \sum q^d \cdot ((d_0 d)! \cdot (e^{\alpha_X} I^X)_{d,H^0})$. Consider the operator (which generalizes the Riemann–Roch operator, see* [Go01]*)*

$$L = \prod_{i=1}^{k} \big(w_i D - (w_i - 1)\big)_{\boldsymbol{w_i}} - q \prod_{i=0}^{l} \big(d_i D + 1\big)_{\boldsymbol{d_i}}.$$

Then $L[\mathcal{I}^X(q)] = 0$.

1.2. **Theorem.** *Let Y be a $\mathbb{Q}$-factorial toric variety and $Y_1, \ldots, Y_k$ be the divisors that correspond to the edges of the fan of Y. Consider a smooth complete intersection X of hypersurfaces $X_1, \ldots, X_l$ that does not intersect the singular locus of Y. Assume that $-K_X > 0$ and $\operatorname{Pic} X = \mathbb{Z}$. Let $i\colon X \to Y$ be the natural embedding. Let ℓ be a nef generator of $H_2(Y)$. For $\beta = d\ell$ put $q^\beta = q^d$. Let $\Lambda \subset H_2(X)$ be the semigroup of algebraic curves as cycles on X.*

1) *I-series of X is the following.*

$$I^X = e^{-\alpha_X q} \sum_{\beta \in \Lambda} q^\beta \cdot i^* \left(\frac{\prod_{a=1}^{l} \big(X_a + 1\big)_{\boldsymbol{\beta \cdot X_a}}}{\prod_{a=1}^{k} \big(Y_a + 1\big)_{\boldsymbol{\beta \cdot Y_a}}} \right),$$

where $\alpha_X = 0$ if the index of X is greater than 1, and $\alpha_X = \prod_{a=1}^{l} (\ell \cdot X_a)!/\prod_{a=1}^{k} (\ell \cdot Y_k)!$ if the index is 1.

2) *Let d_i $(1 \le i \le l)$ be the degrees of the hypersurfaces X_i (with respect to ℓ), w_i $(1 \le i \le k)$ be the degrees of divisors that correspond to the edges of the fan of Y, and d_0 be the index of X.Let $\mathcal{I}^X(q) = \sum q^d \cdot ((d_0 d)! \cdot (e^{\alpha_X} I^X)_{d,H^0})$. Consider the operator (which generalizes the Riemann–Roch operator, see* [Go01]*)*

$$L = \prod_{i=1}^{k} \big(w_i D - (w_i - 1)\big)_{\boldsymbol{w_i}} - q \prod_{i=0}^{l} \big(d_i D + 1\big)_{\boldsymbol{d_i}}.$$

Then $L[\mathcal{I}^X(q)] = 0$.

Remark 1. By Lefschetz Theorem (see [Do82], Theorem 3.2.4, (i) and Remark 3.2.6) hypotheses of theorems 1.1 and 1.2 hold for complete intersections of dimension greater than 2, which do not intersect the singular locus of ambient variety, if its Picard group is $\mathbb{Z}$ (this holds automatically by hypothesis of theorem 1.1).

2. Preliminaries

Throughout the paper we consider the invariants of *genus zero* (those that correspond to the rational curves).

An axiomatic treatment of *prime invariants* was given by M. Kontsevich and Yu. Manin in [KM94]. *Invariants with descendants* were introduced and constructed in [BM96] and [Beh97].

2.1. **Moduli spaces of curves.**

Consider smooth variety X such that $-K_X \geq 0$.

2.1.1. **Definition**. The curve *is reduced scheme of pure dimension* 1. The genus *of curve* C *is the number* $h^1(\mathcal{O}_C)$.

It is easy to see that the curve is of genus 0 if and only if it is a tree of rational curves.

2.1.2. **Definition**. *The connected curve* C *with* $n \geq 0$ *marked points* $p_1, \ldots, p_n \in C$ *is called* prestable *if it has at most ordinary double points as singularities and* $p_1, \ldots, p_n$ *are distinct smooth points* (*see* [Ma02], *III–2.1*). *The map* $f\colon C \to X$ *of connected curve of genus* 0 *with* n *marked points are called* stable *if* C *is prestable and there are at least three marked or singular points on every contracted component of* C ([Ma02], *V–1.3.2*).

In the other words, a stable map of connected curve is the map that has only finite number of infinitesimal automorphisms.

2.1.3. **Definition**. The family of stable maps (*over the scheme* S) *of curves of genus* 0 *with* n *marked points is the collection* $(\pi\colon \mathcal{C} \to S, p_1, \ldots, p_n, f\colon \mathcal{C} \to X)$, *where* π *is the following map. It is a smooth projective map with* n *sections* $p_1, \ldots, p_n$. *Its geometric fibers* $(\mathcal{C}_s, p_1(s), \ldots, p_n(s))$ *are prestable curves of genus* 0 *with* n *marked points. Finally, the restriction* $f|_{\mathcal{C}_s}$ *on each fiber is a stable map.*

Two families over S

$$(\pi\colon \mathcal{C} \to S, p_1, \ldots, p_n, f), \quad (\pi'\colon \mathcal{C}' \to S, p'_1, \ldots, p'_n, f')$$

are called isomorphic *if there is an isomorphism* $\tau\colon \mathcal{C} \to \mathcal{C}'$ *such that* $\pi = \pi' \circ \tau$, $p'_i = \tau \circ p_i$, $f = f' \circ \tau$.

2.1.4. Let $\beta \in H_2^+(X)$. Consider the following (contravariant) functor $\overline{\mathcal{M}}_n(X, \beta)$ from the category of (complex algebraic) schemes to the category of sets. Let $\overline{\mathcal{M}}_n(X, \beta)(S)$ be the set of isomorphism classes of families of stable maps of genus 0 curves with n marked points $(\pi\colon \mathcal{C} \to S, p_1, \ldots, p_n, f)$ such that $f_*([\mathcal{C}_s]) = \beta$, where $[C_s]$ is the fundamental class of C_s.

2.1.5. **Definition**. The moduli space of stable maps *of genus* 0 *curves of class* $\beta \in H_2^+(X)$ *with* n *marked points to* X *is the Deligne–Mumford stack* (*see* [Ma02], *V–5.5*) *which is the coarse moduli space* (*see* [HM98], *Definition* 1.3) *that represents* $\overline{\mathcal{M}}_n(X, \beta)$. *This space is denoted by* $\bar{M}_n(X, \beta)$.

2.2. **Gromov–Witten invariants and I-series.** The definition of Gromov–Witten invariants is given in terms of intersection theory on smooth stacks $\bar{M}_n(X,\beta)$. It exists because locally such stack is a quotient of smooth variety by a finite group (see [Vi89]). However such stacks may have unexpected dimension. To use the products of cohomological cycles on them one should introduce *the virtual fundamental class* $[\bar{M}_n(X,\beta)]^{\mathrm{virt}}$ of virtual dimension $\mathrm{vdim}\,\bar{M}_n(X,\beta) = \dim X - \deg_{K_X}\beta + n - 3$ (see its construction in [Ma02], VI–1.1). In many cases (for instance, for homogenous spaces) virtual fundamental class coincides with the usual one.

Consider the map $ev_i\colon \bar{M}_n(X,\beta) \to X$, $ev_i(C;p_1,\ldots,p_n,f) = f(p_i)$. Let $\pi_{n+1}\colon \bar{M}_{n+1}(X,\beta) \to \bar{M}_n(X,\beta)$ be the forgetful map at the point p_{n+1}. It contracts unstable components after forgetting p_{n+1}. Let $\sigma_i\colon \bar{M}_n(X,\beta) \to \bar{M}_{n+1}(X,\beta)$ be the section that are correspond to the marked point p_i constructed as follows. The image of the curve $(C;p_1,\ldots,p_n,f)$ under the map σ_i is the curve $(C';p_1,\ldots,p_{n+1},f')$. Here $C' = C\bigcup C_0$, $C_0 \simeq \mathbb{P}^1$, and C_0 and C intersect at the (non-marked on C') point p_i. The points p_{n+1} and new p_i lie on C_0. The map f' contracts C_0 and $f'|_C = f$.



Let $L_i = \sigma_i^*\omega_{\pi_{n+1}}$, where $\omega_{\pi_{n+1}}$ is the relative dualizing sheaf π_{n+1}. The fiber L_i over the point $(C;p_1,\ldots,p_n,f)$ is $T^*_{p_i}C$.

2.2.1. **Definition** [see [Ma02], VI–2.1]. A cotangent line class *is the class*

$$\psi_i = c_1(L_i) \in H^2(\bar{M}_n(X,\beta)).$$

2.2.2. **Definition** [see [Ma02], VI–2.1]. *Consider the cohomological classes* $\gamma_1,\ldots,\gamma_n \in H^*(X)$. *Let* $a_1,\ldots,a_n$ *be the non-negative integers and* $\beta \in H_2(X)$. The Gromov–Witten invariant with descendants *that correspond to these classes is*

$$\langle\tau_{a_1}\gamma_1,\ldots,\tau_{a_n}\gamma_n\rangle_\beta = \psi_1^{a_1}ev_1^*(\gamma_1)\cdot\ldots\cdot\psi_n^{a_n}ev_n^*(\gamma_n)\cdot[\bar{M}_n(X,\beta)]^{\mathrm{virt}}$$

if $\sum\mathrm{codim}\,\gamma_i + \sum a_i = \mathrm{vdim}\,\bar{M}_n(X,\beta)$ *and* 0 *otherwise. The invariants with* $a_i = 0$ *(for each* i*) are called* prime. *They are equal to the expected numbers of rational curves of the class* β *on* X*, which intersect the dual cycles to* $\gamma_1,\ldots,\gamma_n$ *(divided by the product of degrees of divisors from* $\gamma_1,\ldots,\gamma_n$*). We omit the symbols* τ_0.

2.2.3. **Definition** [see [Ga00]]. *Let* $\mu_1, \dots, \mu_N$ *be the basis of* $H^*(X)$, $\check{\mu}_1, \dots, \check{\mu}_N$ *be the dual basis, and* $\gamma_1, \dots, \gamma_k$ *be the basis of* $H_2(X)$. *Each curve* β *is of the form* $\beta = \sum \beta_i \gamma_i$. *Consider the ring* $B = \mathbb{C}[[q]]$, $q = (q_1, \dots, q_k)$. *Put* $q^\beta = \prod q_i^{\beta_i}$. *Then* the I-series *for* X *is the following.*

$$I^X = \sum_{\beta \geq 0} I^X_\beta \cdot q^\beta \in B \otimes H^*(X), \qquad I^X_\beta = \sum_{i,j \geq 0} \langle \tau_i \mu_j \rangle_\beta \check{\mu}_j$$

(*we use the same symbol for* $\gamma \in H^*(X)$ *and* $1 \otimes \gamma \in B \otimes H^*(X)$).

Fundamental term *of* I*-series is the series*

$$I^X_{H^0} = \sum_{\beta \geq 0} \langle \tau_{(-K_X)\cdot\beta - 2} \check{\mathbf{1}} \rangle_\beta \cdot q^\beta,$$

where $\check{\mathbf{1}}$ *is the class dual to the class of unity in the cohomology.*

In the case $k = 1$ and arbitrary series $I \in B \otimes H^*(X)$ put $I = \sum_{q \geq 0} I_d \cdot q^d$.

2.2.4. **Lemma** [[Ga99], Lemma 5.5 or proof of Lemma 1 in [LP01]]. *Let* $Y \subset X$ *be a complete intersection and* $\varphi\colon H^*(X) \to H^*(Y)$ *be the restriction homomorphism. Let* $\widetilde{\gamma}_1 \in \varphi(H^*(X))^\perp$ *and* $\gamma_2, \dots, \gamma_l \in \varphi(H^*(X))$. *Then for each* $\beta \in \varphi(H_2(X)) \subset H_2(Y)$ *the Gromov–Witten invariant on* Y *of the form*

$$\langle \tau_{d_1} \widetilde{\gamma}_1, \tau_{d_2} \gamma_2, \dots \tau_{d_l} \gamma_l \rangle_\beta$$

vanishes.

2.2.5. **Definition**. *Consider a complete intersection* $Y \subset X$. *Let* $\varphi\colon H^*(X) \to H^*(Y)$ *be the restriction homomorphism and* $R = \varphi(H^*(X)) \subset H^*(Y)$. *Let* $\mu_1, \dots, \mu_N$ *be the basis of* R, $\check{\mu}_1, \dots, \check{\mu}_N \in R$ *be the dual basis, and* $\gamma_1, \dots, \gamma_k$ *be the basis of* $H_2(Y)$. *Each curve* β *is of the form* $\beta = \sum \beta_i \gamma_i$. *Consider the ring* $B = \mathbb{C}[[q]]$, $q = (q_1, \dots, q_k)$. *Put* $q^\beta = \prod q_i^{\beta_i}$. *Then* restricted I-series *of* Y *is the following.*

$$\widetilde{I}^Y = \sum_{\beta \geq 0} \widetilde{I}^Y_\beta \cdot q^\beta \in B \otimes R \subset B \otimes H^*(Y), \qquad \widetilde{I}^Y_\beta = \sum_{i,j \geq 0} \langle \tau_i \mu_j \rangle_\beta \check{\mu}_j.$$

The invariants of complete intersection that correspond to cohomological classes restricted from the ambient variety are called restricted.

2.2.6. **Remark**. For a complete intersection $Y \subset X$ of dimension at least 3 these series coincide, i. e. $I^Y = \widetilde{I}^Y$. Indeed, $H_2(Y) \simeq H_2(X)$ and by lemma 2.2.4 and the divisor axiom 2.3.2 one-pointed invariants for primitive classes vanish.

2.2.7. **Remark**. Two- and more- pointed invariants for primitive classes of middle dimension may be non-zero. For instance, for two primitive classes α and β, a hypersurface section H and a line ℓ on cubic threefold

$$\langle \alpha, \beta, H^2 \rangle_\ell = -\alpha \cdot \beta$$

(see [Bea95], Proposition 1).

2.3. Relations between multipointed invariants.

2.3.1. In this paragraph we discuss relations between one-pointed Gromov–Witten invariants with descendants (i. e. coefficients of I-series) and multipointed prime ones (in particular, two-pointed ones, which are important for the following).

One-pointed invariants with descendants may be expressed in terms of multipointed ones by the following.

2.3.2. **Divisor axiom for invariants with descendants** [[KM98], 1.5.2]. *Let Y be smooth projective variety, $\gamma_1, \ldots, \gamma_n \in H^*(Y)$ and $\gamma_0 \in H^2(Y)$ be a divisor. Then*

$$\langle \gamma_0, \tau_{d_1}\gamma_1, \ldots, \tau_{d_n}\gamma_n \rangle_\beta = (\gamma_0 \cdot \beta)\langle \tau_{d_1}\gamma_1, \ldots, \tau_{d_n}\gamma_n \rangle_\beta + \sum_{k, d_k \geq 1} \langle \tau_{d_1}\gamma_1, \ldots, \tau_{d_k-1}(\gamma_0 \cdot \gamma_k), \ldots, \tau_{d_n}\gamma_n \rangle_\beta.$$

The following formula (with divisor axiom) enables one to express recursively three-pointed invariants with descendants in terms of prime ones.

2.3.3. **Theorem** [topological recursion relations, [Ma02], VI–6.2.1]. *Let Y be a smooth projective variety, $\{\Delta^i\}$ and $\{\Delta_i\}$, $i = 1, \ldots, N$, be dual bases of $H^*(Y)$. Then*

$$\langle \tau_{d_1}\gamma_1, \tau_{d_2}\gamma_2, \tau_{d_3}\gamma_3 \rangle_\beta = \sum_{a, \beta_1+\beta_2=\beta} \langle \tau_{d_1-1}\gamma_1, \Delta^a \rangle_{\beta_1} \langle \Delta_a, \tau_{d_2}\gamma_2, \tau_{d_3}\gamma_3 \rangle_{\beta_2}$$

(*the sum is taken over all a, β_1, β_2 such that the expression in it is well-defined, i. e. $a = 1, \ldots, N$, $\beta_1, \beta_2 \geq 0$*).

2.3.4. Using this formulas, we can express one-pointed invariants in terms of prime two-pointed ones (see [Pr04], Proposition 5.2). In fact, in many cases these expressions are invertible. More particular, the inverse expressions can be found recursively by the divisor axiom and the following theorem (see also [BK00], 5.2 and [Pr04]).

2.3.5. **Theorem** [Lee, Pandharipande, [LP01], Theorem 2, i]. *Let Y be a smooth projective variety. Consider the self-dual ring $R \subset H^*(Y)$ generated by Picard group $\operatorname{Pic} Y$ such that for any $\mu_1, \ldots, \mu_n \in R$ and $\nu \in R^\perp$*

$$\langle \tau^{a_1}\mu_1, \ldots, \tau^{a_n}\mu_n, \tau^a\nu \rangle_\beta = 0.$$

Let $\gamma_1, \ldots, \gamma_n \in R$, $H \in \operatorname{Pic} Y$, Δ^i and Δ_i be the dual bases of $H^*(Y)$. Then one can algebraically express Gromov–Witten invariant

$$\langle \tau^{k_1}\gamma_1, \ldots, \tau^{k_n}\gamma_n \rangle_\beta$$

in terms of one-pointed Gromov–Witten invariants with descendants using the following expressions

$$\langle \tau_{k_1}\gamma_1, \ldots, \tau_{k_n}H \cdot \gamma_n \rangle_\beta = \langle \tau_{k_1}H \cdot \gamma_1, \ldots, \tau_{k_n}\gamma_n \rangle_\beta + \beta H \cdot \langle \tau_{k_1+1}\gamma_1, \ldots, \tau_{k_n}\gamma_n \rangle_\beta - \sum_{\beta_1+\beta_2=\beta} \beta_1 H \cdot \sum_{S^1 \cup S^n = S,\ a} \langle \tau_{k_{s_1^1}}\gamma_{s_1^1}, \ldots, \tau_{k_{s_a^1}}\gamma_{s_a^1}, \Delta^a \rangle_{\beta_1} \cdot \langle \Delta_a, \tau_{k_{s_1^n}}\gamma_{s_1^n}, \ldots, \tau_{k_{s_b^n}}\gamma_{s_b^n} \rangle_{\beta_2}$$

and

$$\langle \tau_{k_1}\gamma_1, \ldots, \tau_{k_n+1}\gamma_n \rangle_\beta = -\langle \tau_{k_1+1}\gamma_1, \ldots, \tau_{k_n}\gamma_n \rangle_\beta + \sum_{\substack{\beta_1+\beta_2=\beta,\\ S^1 \cup S^n = S,\ a}} \langle \tau_{k_{s_1^1}}\gamma_{s_1^1}, \ldots, \tau_{k_{s_a^1}}\gamma_{s_a^1}, \Delta^a \rangle_{\beta_1} \cdot \langle \Delta_a, \tau_{k_{s_1^n}}\gamma_{s_1^n}, \ldots, \tau_{k_{s_b^n}}\gamma_{s_b^n} \rangle_{\beta_2},$$

where $H \in \operatorname{Pic} X \cap R$ *and the latter summations are taken over partitions* $S^1 = \{s_1^1, \ldots, s_a^1\}$ *and* $S^n = \{s_1^n, \ldots, s_b^n\}$ *of the set* $S = \{1, \ldots, n\}$ *such that* $1 \in S^1$ *and* $n \in S^n$.

2.3.6. The hypothesis of this theorem holds for subring of restricted cohomologies of complete intersection in toric variety (see [Fu93], 5.2) or for the subring of the cohomology ring of Fano threefold generated by Picard group. See expressions for Fano threefolds with Picard group $\mathbb{Z}$ in [Pr04].

Consequently, theorems 1.1 and 1.2 enable us to find the restricted quantum cohomology ring of smooth complete intersections in weighted projective spaces and singular toric varieties.

2.4. **Toric varieties.** The definition and the main properties of toric varieties see in [Da78] or in [Fu93]. Just remind that toric variety is a variety with action of torus $T \simeq (\mathbb{C}^*)^n$ such that one of its orbits is a Zariski open set. Toric variety is determined by its *fan*, i. e. some collection of cones with vertices in the points of lattice that is dual to the lattice of torus characters. Moreover, algebraic-geometric properties of toric variety can be formulated in terms of properties of this fan.

Remind some of them.

2.4.1. Every cone of the fan $\sigma \subset N = \mathbb{R}^n$ of dimension r corresponds to the orbit of the torus of dimension $n - r$ (here n is the dimension of toric variety). Thus, each edge (one-dimensional cone) correspond to the (equivariant) Weil divisor[1]. The divisors which correspond to the edges of the fan generate divisor class group. A Weil divisor $D = \sum d_i M_i$, where M_i corresponds to edges, is Cartier if for each cone of the fan σ there exist a vector n_σ such that $\langle n_\sigma, m_i \rangle = d_i$ where m_i is the primitive elements of the edges of this cone. If such vector is the same for all cones, then the divisor is principal. Hence if the toric variety is n-dimensional and the number of the edges is k, then the rank of the divisor class group is $n - k$.

2.4.2. **Definition**. *The variety is called* $\mathbb{Q}$-factorial *if for each Weil divisor* D *there exist some integer* k *such that* kD *is a Cartier divisor.*

In particular, there exist an intersection theory for Weil divisors on the $\mathbb{Q}$-factorial variety.

2.4.3. Toric variety is $\mathbb{Q}$-factorial if and only if any cone of the fan, which corresponds to this variety, is simplicial. In this case Picard group is generated (over $\mathbb{Q}$) by divisors, which correspond to the edges of the fan.

2.4.4. Consider a weighted projective space $\mathbb{P} = \mathbb{P}(w_0, \ldots, w_l)$. The fan which correspond to it is generated by integer vectors $m_0, \ldots, m_l \in \mathbb{R}^l$ such that $\sum w_i m_i = 0$. If $w_0 = 1$, then one can put $m_0 = (-w_1, \ldots, -w_l)$, $m_i = e_i$, where e_i is a basis of $\mathbb{R}^l$. The collection $\{m_i\}$ corresponds to the collection of standard divisors–strata $\{D_i \in |w_i H|\}$.

---

[1] That is, let $\Sigma \in N = \mathbb{Z}^n$ be a fan of the toric variety X_Σ and let $\sigma \in \Sigma$ be any cone. Let M be a lattice dual to N with respect to some non-degenerate pairing $\langle \cdot, \cdot \rangle$ and $\sigma^\vee$ be a dual cone for σ (i. e. $\sigma^\vee = \{l \in M | \forall k \in \sigma \ \ \langle l, k \rangle \geq 0\}$). Let $U_\sigma = \operatorname{Spec} \mathbb{C}[\sigma^\vee]$ corresponds to σ. The variety X_σ is obtained from the affine varieties U_σ, $\sigma \in \Sigma$, by gluing together U_σ and U_τ along $U_{\sigma \cap \tau}$, $\sigma, \tau \in \Sigma$. Thus, if $l \subset \sigma \in \Sigma$ is an edge of the fan, then the divisor that is correspond to l restricted on U_σ as $U_l \subset U_\sigma$.

2.4.5. A toric variety is *smooth* if for any cone σ in the fan that correspond to this variety the subgroup $\sigma \cap \mathbb{Z}^n$ is generated by the subset of the basis of the lattice $m_1^\sigma, \ldots, m_k^\sigma$. Adding the edge $a = a_1 m_1^\sigma + \ldots + a_k m_k^\sigma$, $a_i \in \mathbb{Q}$ to the cone (and connection it with "neighboring" faces) corresponds to weighted blow-up (along subvariety which correspond to σ) with weights $1/r \cdot (\alpha_1, \ldots, \alpha_k)$, where $\alpha_i \in \mathbb{Z}$ and $a_i = \alpha_i / r$. Consecutively adding edges to the fan in this way we can get toric resolution of a toric variety.

2.4.6. So, singular locus of weighted projective space $\mathbb{P} = \mathbb{P}(w_0, \ldots, w_l)$ is the union of strata given by $x_{i_1} = \ldots = x_{i_j} = 0$, where x_{i_j} is the coordinate of weight w_{i_j} and $\{i_1, \ldots, i_j\}$ is the maximal set of indices such that greatest common factor of the others is greater than 1.

2.5. **Givental's Theorem** [[Gi97], Theorem 0.1]. *Let X be a smooth toric variety and Y be a smooth complete intersection in it with positive anticanonical class. Let $X_1, \ldots, X_k$ be the divisors which correspond to the edges of a fan of X and let Y be given by divisors $Y_1, \ldots, Y_r$. Let $\Lambda \subset H_2(X)$ be the semigroup of algebraic curves as cycles on Y, and $i \colon Y \to X$ be a natural embedding. Then*

$$I^Y = e^{h(q)} \sum_{\beta \in \Lambda} q^\beta \cdot i^* \left(\frac{\prod_{a=1}^r \left[Y_a\right]_{\boldsymbol{\beta \cdot Y_a + 1}}}{\prod_{a=1}^r \left[Y_a\right]_{\mathbf{1}}} \frac{\prod_{a=1}^k \left[X_a\right]_{\mathbf{1}}}{\prod_{a=1}^k \left[X_a\right]_{\boldsymbol{\beta \cdot X_a + 1}}} \right),$$

where $h(q)$ is a polynomial supported by curves, whose intersection with anticanonical class is 1 (i. e. $h(q) = \sum_\beta h_\beta q^\beta$, where $\beta \cdot (-K_Y) = 1$).

2.6. **Motivation.** Consider a smooth Fano threefold X with Picard group $\mathbb{Z}$. Put $K = -K_X$.

2.6.1. **Definition** [[Go02], 1.7, 1.10]. A counting matrix *is the matrix of Gromov-Witten invariants of X, namely the following matrix $A \in \mathrm{Mat}(4 \times 4)$.*

$$A = \begin{bmatrix} a_{00} & a_{01} & a_{02} & a_{03} \\ 1 & a_{11} & a_{12} & a_{13} \\ 0 & 1 & a_{22} & a_{23} \\ 0 & 0 & 1 & a_{33} \end{bmatrix}.$$

Numeration of rows and columns starts from 0 and the elements are given by

$$a_{ij} = \frac{\langle K^{3-i}, K^j, K \rangle_{j-i+1}}{\deg X} = \frac{j-i+1}{\deg X} \cdot \langle K^{3-i}, K^j \rangle_{j-i+1}$$

(*the degree is taken with respect to the anticanonical class*).

2.6.2. It is easy to see that the matrix A is symmetric with respect to the secondary diagonal: $a_{ij} = a_{3-j,3-i}$. By definition, $a_{ij} = 0$ if $j - i + 1 < 0$. If $j - i + 1 = 0$, then $a_{ij} = 1$, because it is just a number of intersection points of K^{3-i}, K^j, and K, which is $\deg X$; $a_{00} = a_{33} = 0$. For the other coefficients a_{ij}'s are "expected" numbers of rational curves of degree $j - i + 1$ passing through K^{3-i} and K^j, multiplied by $\frac{j-i+1}{\deg X}$. The only exception is the following: by divisor axiom

$$a_{01} = 2 \cdot (2 \cdot \mathit{ind}\,(X) \cdot [\textit{the number of conics passing through the general point}]).$$

2.6.3. Consider the following Fano threefolds (see [Is77], [Is78], [Is79], [Is88], [IP99], [Mu92]).

1: The variety V_1 of anticanonical degree 8 (a double covering of the cone over Veronese surface branched over a smooth cubic).

2: The variety V_2 of anticanonical degree 16 (a double covering of $\mathbb{P}^3$ branched over a smooth quartic).

3: The variety V_2' of anticanonical degree 2 (a double covering of $\mathbb{P}^3$ branched over a smooth sextic).

2.6.4. **Proposition**. *The varieties V_1, V_2, and V_2' are of the following form.*

1: *Any variety of type V_1 is isomorphic to a smooth hypersurface of degree 6 in $\mathbb{P}(1,1,1,2,3)$.*

2: *Any variety of type V_2 is isomorphic to a smooth hypersurface of degree 4 in $\mathbb{P}(1,1,1,1,2)$.*

3: *Any variety of type V_2' is isomorphic to a smooth hypersurface of degree 6 in $\mathbb{P}(1,1,1,1,3)$.*

2.6.5. **Proof**. Double covering $\mathbb{P}(w_0,\ldots,w_n)$ branched over a divisor given by function $f_k(x_0,\ldots,x_n)$ of degree k may be given by function $x_{n+1}^2 = f_k$ in $\mathbb{P}(w_0,\ldots,w_n,k/2)$. The variables x_i here have the weights w_i and the variable x_{n+1} has the weight $k/2$. $\square$

2.6.6. **Theorem**. *The counting matrices for V_1, V_2, and V_2' are as follows.*

1: *For V_1*

$$M(V_1) = \begin{bmatrix} 0 & 240 & 0 & 57600 \\ 1 & 0 & 1248 & 0 \\ 0 & 1 & 0 & 240 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

2: *For V_2*

$$M(V_2) = \begin{bmatrix} 0 & 48 & 0 & 2304 \\ 1 & 0 & 160 & 0 \\ 0 & 1 & 0 & 48 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

3: *For V_2'*

$$M(V_2') = \begin{bmatrix} 0 & 137520 & 119681240 & 21690374400 \\ 1 & 624 & 650016 & 119681240 \\ 0 & 1 & 624 & 137520 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

2.6.7. **Proof.** By proposition 2.6.4 these varieties are hypersurfaces in weighted projective spaces. Find their one-pointed invariants using theorem 1.1. Apply theorem 2.3.5 for the subring of cohomology ring generated by the class dual to the class of hyperplane section and find prime two-pointed invariants. They are the coefficients of counting matrices. $\square$

3. Proofs of the main theorems

3.1. Proof of theorems 1.1 and 1.2.

The divisors in $\mathbb{P}$ that correspond to the edges of its fan are $w_1H, \ldots, w_kH$. Thus, theorem 1.1 follows from theorem 1.2.

Prove theorem 1.2.

1) Let Σ be a fan of Y. Consider the sequence of simplicial fans $\Sigma = \Sigma_0, \Sigma_1, \ldots, \Sigma_r$ such that

- i) The fan Σ_{i+1} is a result of the following procedure applied to Σ_i. Pick a cone whose edges are not part of a basis of the integer lattice containing this fan. Add an edge, which lies inside the cone. Add the faces which contain this edge and "neighboring" edges (i. e. replace all cones which contain the edge we add by all linear spans of this edge and faces of this cone that does not contain the edge);
- ii) The fan Σ_r corresponds to a smooth variety.

Such sequence exists by [Da78], 8.1–8.3.

This way we get a toric resolution of singularities $f\colon \widetilde{Y} = X_{\Sigma_r} \to Y$ (where X_{Σ_r} is the toric variety that correspond to the fan Σ_r, see remark 1). The exceptional set of this resolution is the union of divisors $E_1, \ldots, E_r$ which correspond to the edges we add. Let $W_1, \ldots, W_k$ be divisors that correspond to the edges of a fan of Y and $\widetilde{W}_1, \ldots, \widetilde{W}_k$ be their strict transforms. Let $\widetilde{X}_1, \ldots, \widetilde{X}_l$ be strict transforms of $X_1, \ldots, X_l$, $\widetilde{X} = \widetilde{X}_1 \cap \ldots \cap \widetilde{X}_l$, and $j\colon \widetilde{X} \to \widetilde{Y}$ be the natural embedding. Obviously, there exists an isomorphism g such that the diagram

$$\begin{tikzcd}
\widetilde{X} \arrow[r, "j"] \arrow[d, "g"'] & \widetilde{Y} \arrow[d, "f"] \\
X \arrow[r, "i"] & Y
\end{tikzcd}$$

is commutative, because f is an isomorphism between $\widetilde{Y} \setminus (\cup_{i=1}^{r} E_i)$ and $Y \setminus \operatorname{Sing} Y$.

Consider any toric variety V associated to a complete simplicial fan. Then the canonical map of group of algebraic cycles modulo rational equivalence with rational coefficients to homology group with rational coefficients

$$A_*(V)_{\mathbb{Q}} \to H_*(V, \mathbb{Q})$$

is an isomorphism (see [Da78], 10.9). So, we can extend an intersection theory to the homology group with rational coefficients.

The map $f_*\colon H_2(\widetilde{Y}, \mathbb{Q}) \to H_2(Y, \mathbb{Q})$ is surjective. Let $\lambda \subset X \subset Y$ be an effective curve such that $H_2(X) = \mathbb{Z}\lambda$ and $\widetilde{\lambda} = f^{-1}(\lambda)$. Put $K = \ker f_*$. Then $H_2(\widetilde{Y}, \mathbb{Q}) = \mathbb{Q}\widetilde{\lambda} + K$.

The cycle of any curve $\boldsymbol{\beta}$ which lies on $\widetilde{X}$ equals $\beta_0 \cdot \widetilde{\lambda}$, where $\beta_0 \in \mathbb{Z}$ and $\beta_0 \geq 0$, because representatives of elements of K lie on exceptional divisors, which do not intersect $\widetilde{X}$. So, if $\widetilde{\Lambda}$ is a semigroup of algebraic curves on $\widetilde{X}$, then $\widetilde{\Lambda} = \mathbb{Z}_{\geq 0}\widetilde{\lambda}$.

The multiplicity of intersection of divisor and a curve depends only on the neighborhood of this curve and f is an isomorphism in a neighborhood of $\widetilde{X}$, so $\widetilde{\lambda} \cdot \widetilde{X}_i = \lambda \cdot X_i$ and $\widetilde{\lambda} \cdot \widetilde{W}_i = \lambda \cdot W_i$. Thus, for any curve $\boldsymbol{\beta} = \beta_0 \cdot \widetilde{\lambda}$ on $\widetilde{X}$ we have $\boldsymbol{\beta} \cdot \widetilde{W}_i = \beta_0 \cdot (\lambda \cdot W_i)$, $\boldsymbol{\beta} \cdot \widetilde{X}_i = \beta_0 \cdot (\lambda \cdot X_i)$, and $\boldsymbol{\beta} \cdot E_i = 0$.

Divisors that correspond to the edges of a fan of $\widetilde{Y}$ are $\widetilde{W}_1, \ldots, \widetilde{W}_k, E_1, \ldots, E_r$. But the factors in the expression for I-series for $\widetilde{X}$ from Theorem 2.5 which include the divisors E_i may be cancelled, so we may omit them. Consider rings $H^*(X)[[q]]$ and $H^*(\widetilde{X})[[\widetilde{q}]]$

and an isomorphism between them which sends q to $\widetilde{q}$ and acts on the coefficients as f^*. Let $j\colon \widetilde{X} \to \widetilde{Y}$ be a natural embedding. Then

$$I^{\widetilde{X}} = e^{\widetilde{h}(\widetilde{q})} \sum_{\beta=\beta_0\widetilde{\lambda}\in\widetilde{\Lambda}} \widetilde{q}^{\beta_0} \cdot j^* \left(\frac{\prod_{a=1}^{l} \big[\widetilde{X}_a\big]_{\boldsymbol{\beta\cdot \widetilde{X}_a+1}}}{\prod_{a=1}^{l} \big[\widetilde{X}_a\big]_{\mathbf{1}}} \frac{\prod_{a=1}^{k} \big[\widetilde{W}_a\big]_{\mathbf{1}}}{\prod_{a=1}^{k} \big[\widetilde{W}_a\big]_{\boldsymbol{\beta\cdot \widetilde{W}_a+1}}} \right) =$$

$$j^*f^* \left(e^{h(q)} \sum_{\beta_0\geq 0} q^{\beta_0} \cdot \prod_{a=1}^{l} \big(X_a+1\big)_{\boldsymbol{\beta_0\cdot \mathbf{deg}\, X_a}} \cdot \frac{1}{\prod_{a=1}^{k} \big(\widetilde{W}_a+1\big)_{\boldsymbol{\beta_0\cdot w_a}}} \right) =$$

$$g^* \left(e^{h(q)} \sum_{d=0}^{\infty} q^d \cdot i^* \left(\frac{\prod_{a=1}^{l} \big(X_a+1\big)_{\boldsymbol{d\cdot \mathbf{deg}\, X_a}}}{\prod_{a=1}^{k} \big(\widetilde{W}_a+1\big)_{\boldsymbol{d\cdot w_a}}} \right) \right),$$

where $\widetilde{h}(\widetilde{q})$ is a polynomial from theorem 2.5 and $h(q)$ is its pre-image. The second equality holds because we may choose divisors that intersect X properly, and their strict transforms as representatives (over $\mathbb{Q}$) of divisor classes (because $\operatorname{Pic} Y = \mathbb{Z}$ and every effective divisor on Y is ample, and, therefore, it is equivalent to a rational multiple to a very ample one). The map f after restriction on $\widetilde{X}$ is isomorphism, which induces the isomorphism f^* of cohomology groups such that the image of any divisor under f^* is its strict transform. Finally, g^* is an isomorphism of cohomology rings X and $\widetilde{X}$, so $I^{\widetilde{X}} = g^*(I^X)$.

The only thing that has to be found is $h(q)$. If the index of X is greater than 1, then there are no curves which intersect the anticanonical class by 1, so $h(q)=0$. If the index is 1, then any such curve is a line ℓ (with respect to the anticanonical class of X) and $h(q) = -\alpha_X q$. Besides, there are no lines on X passing through the general point, i. e. $\langle H^{\dim X}\rangle_\ell = 0$. This means that the coefficient $e^{-\alpha_X q} \cdot I^X_\ell$ at H^0 vanishes, so

$$\alpha_X = \frac{\prod_{a=1}^{l}(\ell\cdot X_a)!}{\prod_{a=1}^{k}(\ell\cdot Y_k)!} = \frac{\prod_{a=1}^{l}(\deg X_a)!}{\prod_{a=1}^{k} w_a!}.$$

2) It is easy to see that

$$\mathcal{I}^X(q)_{H^0} = \sum_{n=0}^{\infty} \frac{\prod_{i=0}^{l}(d_i\cdot n)!}{\prod_{j=1}^{k}(w_j\cdot n)!} q^n.$$

Consider the solution of the equation $L[\mathcal{J}^X(q)] = 0$ as a series $\mathcal{J}^X = \sum_{n\geq 0} a_n q^n$. Evidently, $D[q^n] = nq^n$. Hence

$$(\prod_{i=1}^{k} \big(w_iD-(w_i-1)\big)_{\boldsymbol{w_i}})[q^n] = (\prod_{i=1}^{k} \big(w_in-(w_i-1)\big)_{\boldsymbol{w_i}})\cdot q^n.$$

Analogously,

$$(q\prod_{i=0}^{l} \big(d_iD+1\big)_{\boldsymbol{d_i}})[q^{n-1}] = (\prod_{i=0}^{l} \big(d_i(n-1)+1\big)_{\boldsymbol{d_i}})\cdot q^n.$$

The differential equation on $\mathcal{J}^X$ translates into recursive equations on the coefficients a_n using these equalities. Namely,

$$a_n = \frac{\prod_{i=0}^{l} \big(d_i(n-1)+1\big)_{\boldsymbol{d_i}}}{\prod_{i=1}^{k} \big(w_i n-(w_i-1)\big)_{\boldsymbol{w_i}}} \cdot a_{n-1}.$$

Put $a_0 = 1$. By induction we get

$$a_n = \sum \frac{\prod_{i=0}^{l}(d_i \cdot n)!}{\prod_{j=1}^{k}(w_j \cdot n)!} q^n.$$

□

3.1.1. **Remark**. It is easy to see that the Riemann–Roch operator is divisible by $D^{\dim Y - \dim X}$ from the left. Let $L = D^{\dim Y - \dim X}\widetilde{L}$. Then $\widetilde{L}\mathcal{I}^X = 0$.

3.2. **Comments and examples**.

3.2.1. **Remarks.**

i: In order to check that a complete intersection $X = X_1 \cup \ldots \cup X_l$ of general hypersurfaces of degrees $d_1, \ldots, d_l$ in a weighted projective space $\mathbb{P} = \mathbb{P}(w_1, \ldots w_k)$ does not intersect the singular locus of $\mathbb{P}$ one can use the following necessary (but not sufficient) condition.

The number of Cartier divisors among X_i is greater than the dimension of the singular locus of $\mathbb{P}$.

This means that

(*a number of d_i's that is divisible on each of a_i's*) $>$ (*the maximal number of weights w_i such that greatest common factor of the others is greater than 1*)

(see 2.4.6).

For an arbitrary toric variety the condition reformulates as follows. The number of movable divisors (i. e. those that we can move from any point) is greater than codimension of maximal cone whose edges are not subset of basis.

ii: We assume that the Picard number of the ambient variety is 1 just for simplicity. In cases of higher Picard number theorems 1.1 and 1.2 may be formulated and proved in a similar manner. We consider a multivariable $q = (q_1, \ldots, q_m)$ instead of the variable q (here $m = \operatorname{rk} \operatorname{Pic} X$) a multidegree of a curve (with respect to the generators of Picard group) instead of a degree and so on. The only difference in this case is the expression for correction term $e^{-\alpha_X q}$.

iii: If we lift the hypothesis $\operatorname{Pic} X = \mathbb{Z}$ (or $\dim X \geq 3$ in remark 1), then theorems 1.1 and 1.2 give *the restricted* I-series (see remark 2.2.6).

iiii: The main theorems can be generalized to Calabi–Yau varieties. Their proofs differ from proofs of theorems 1.1 and 1.2 only in formulas for correction term (see [Gi97]).

3.2.2. Examples.

i: Consider a general hypersurface $X^k \subset \mathbb{P}^n$. Let $k < n$. Then, by theorem 1.1,

$$I^X = \sum_{d=0}^{\infty} q^d \frac{\big(kH+1\big)_{\boldsymbol{kd+1}}}{(\big(H+1\big)_{\boldsymbol{d+1}})^{n+1}}.$$

This coincides with the formula for I-series for a hypersurface in projective space from [Gi96].

If $k = n$, then, by theorem 1.1,

$$I^X = e^{-k!q} \sum_{d=0}^{\infty} q^d \frac{\big(kH+1\big)_{\boldsymbol{kd+1}}}{(\big(H+1\big)_{\boldsymbol{d+1}})^{n+1}}.$$

This coincides with the formula 4.2.2 from [Pr04].

ii: For projective space ($w_i = 1$) theorem 1.1 is Givental's Theorem for complete intersections in projective space.

4. APPENDIX: GOLYSHEV'S CONJECTURE

4.1. Consider *a family* of counting matrices $A^\lambda = A + \lambda \mathbf{1}$, where $\mathbf{1}$ is the identity matrix (see 2.6.1 for the definition of counting matrix). Consider the one-dimensional torus $\mathbb{G}_m = \operatorname{Spec} \mathbb{C}[z, z^{-1}]$ and the differential operator $D = z\frac{\partial}{\partial z}$. To construct the family of matrices M^λ, put its elements m_{kl}^λ as follows

$$m_{kl}^\lambda = \begin{cases} 0, & \text{if } k > l+1, \\ 1, & \text{if } k = l+1, \\ a_{kl}^\lambda \cdot (-\frac{\partial}{\partial z})^{l-k+1}, & \text{if } k < l+1. \end{cases}$$

Consider the family of differential operators

$$\widetilde{L}^\lambda = \det\nolimits_{\text{right}}(D\mathbf{1} - M^\lambda),$$

where $\det_{\text{right}}$ means “right determinant”, i. e. the determinant, which is calculated with respect to *the rightmost* column; all minors are calculated in the same way. Divide $\widetilde{L}^\lambda$ on the left by D. We get the family of operators L^λ, so $\widetilde{L}^\lambda = DL^\lambda$.

4.2. **Definition** [[Go02], 1.8]. *The equation of the family* $L^\lambda[\Phi(z)] = 0$ *is called* counting equation $D3$*.*

4.3. **Golyshev's conjecture** [V. Golyshev, [Go02]]. *The solutions of* $D3$ *equations for Fano threefolds with Picard group* $\mathbb{Z}$ *are modular. More precisely, let* X *be such variety,* i_X *be its index, and* $N = \frac{\deg X}{2i_X^2}$*. Then in the family of counting equations for* X *there is one,* $L^{\lambda_X}[\Phi(z)] = 0$*, whose solution is an Eisenstein series of weight* 2 *on* $X_0(N)$*.*

More precise description of counting equations as Picard–Fuchs equations see in [Go02].

4.4. Based on this conjecture, Golyshev gives a list of predictions of counting matrices of Fano threefolds. To check this conjecture one should find all such matrices.

There are 17 families of smooth Fano threefolds with Picard group $\mathbb{Z}$ (V. Iskovskikh, [Is77], [Is78]). For 14 of them counting matrices were found by A. Beauville ([Bea95]), A. Kuznetsov (unpublished), V. Golyshev (unpublished) and the author ([Pr04]). The matrices for 2 other varieties are given by theorem 2.6.6.

All these matrices coincide with ones that were predicted by Golyshev. Thus, theorem 2.6.6 finishes the proof of Golyshev's conjecture.

The author is grateful to I. Cheltsov, S. Galkin, A. Givental, V. Golyshev, M. Kazarian, S. Shadrin, K. Shramov, M. Tsfasman, and F. Zak for comments.

REFERENCES

[AS72] M. Abramowitz, I. Stegun (eds.) *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing*, New York: Dover, 1972.

[Ba93] V. Batyrev, *Quantum Cohomology Rings of Toric Manifolds*, preprint (1993), alg-geom/9310004.

[Bea95] A. Beauville, *Quantum cohomology of complete intersections*, preprint (1995), alg-geom/9501008.

[Beh97] A. Behrend, *Gromov–Witten invariants in algebraic geometry*, Invent. Math., 127(3), 1997, 601–617.

[BK00] A. Bertram, H. Kley, *New recursions for genus-zero Gromov–Witten invariants*, preprint (2000), math.AG/0007082.

[BM96] K. Behrend, Yu. Manin, *Stacks of Stable Maps and Gromov-Witten Invariants*, Duke Math. J. vol. 85, No. 1 (1996) 1–60, preprint (1995), alg-geom/9506023.

[CJ99] A. Collino, M. Jinzenji, *On the Structure of the Small Quantum Cohomology Rings of Projective Hypersurfaces*, Comm. Math. Phys. 206 (1999) 157-183.

[Da78] V. Danilov, *Geometry of toric varieties*, Russ. Math. Surv., 33 (1978), 97-154.

[Do82] I. Dolgachev, Weighted projective varieties, Lect. Notes Math., 1982, V. 956, 34–71.

[Fu93] W. Fulton, *Introduction to toric varieties*, Princeton University Press, Princeton, NJ, 1993.

[Ga99] A. Gathmann, *Relative Gromov-Witten invariants and the mirror formula*, preprint (1999), math.AG/9908054.

[Ga00] A. Gathmann, *Absolute and relative Gromov-Witten invariants of very ample hypersurfaces*, preprint (2000), math.AG/0009190.

[Gi96] A. Givental, *Equivariant Gromov - Witten Invariants*, IMRN (1996) No. 13, 613-663, preprint (1996), alg-geom/9603021.

[Gi97] A. Givental, *A mirror theorem for toric complete intersections*, preprint (1997), alg-geom/9701016.

[Go01] V. Golyshev, *Riemann-Roch variations*, Izv. Math., 2001, 65 (5), 853-881.

[Go02] V. Golyshev, *Geometricity Problem and Modularity of Certain Riemann–Roch Variations*, Dokl. Math, Vol. 66, No. 2, p. 231 (2002).

[Go04] V. Golyshev, *Modularity of equations $D3$ and the Iskovskikh classification*, Dokl. Math, to appear, preprint (2004), math.AG/0405039.

[HM98] J. Harris, I. Morrison, *Moduli of curves*, Graduate Texts in Mathematics, 187. New York, NY: Springer, xiii, 1998.

[Is77] V. Iskovskikh, *Fano 3-folds I*, Math USSR, Izv. 11 (1977), 485-527.

[Is78] V. Iskovskikh, *Fano 3-folds II*, Math USSR, Izv. 11 (1978), 469-506.

[Is79] V. Iskovskikh, *Anticanonical models of three-dimensional algebraic varieties*, J. Soviet Math. **13** (1980), 745-814.

[Is88] V. Iskovskikh, *Lectures on three-dimensional algebraic varieties. Fano varieties. (Lektsii po trekhmernym algebraicheskim mnogoobraziyam. Mnogoobraziya Fano)*, Moscow, Moscow State University (1988).

[IP99] V. Iskovskikh, Yu. Prokhorov, *Fano varieties*, Encyclopaedia Math. Sci. 47 (1999) Springer, Berlin.

[KM94] M. Kontsevich, Yu. Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Comm. Math. Phys. 164 (1994) 525-562.

[KM98] M. Kontsevich, Yu. Manin, *Relations between the correlators of the topological sigma-model coupled to gravity*, Comm. Math. Phys. 196 (1998) 385-398.

[LP01] Y.-P. Lee, R. Pandharipande, *A reconstruction theorem in quantum cohomology and quantum K-theory*, preprint (2001), math.AG/0104084.

[Ma02] Yu. Manin, *Frobenius manifolds, quantum cohomology, and moduli spaces*, Colloquium Publications. American Mathematical Society (AMS). 47. Providence, RI: American Mathematical Society (AMS) (1999).

[Mu92] S. Mukai, *Fano 3-folds*, Lond. Math. Soc. Lect. Note Ser. 179, 255-263 (1992).

[Pa98] R. Pandharipande, *Rational curves on hypersurfaces [after A. Givental]*, Societe Mathematique de France, Asterisque. 252, 307-340, Exp. No.848 (1998).

[Pr04] V. Przyjalkowski, *Gromov–Witten invariants of Fano threefolds of genera* 6 *and* 8, preprint (2004), math.AG/0410327.

[Vi89] A. Vistoli, *Intersection theory on algebraic stacks and their moduli*, Invent. Math. 97 (1989), 613–670.

[W91a] E. Witten, *Two-dimentional gravity and intersection theory on moduli space*, Surveys in Diff. Geom. 1 (1991), 243-310.

[W91b] E. Witten, *Algebraic geometry associated with matrix models of two dimensional gravity*, Topological methods in modern mathematics (Stony Brook, NY,1991), Publish or Perish, Texas (1993), 235-269.

Steklov Institute of Mathematics, 8 Gubkin street, Moscow 119991, Russia
E-mail address: `victorprz@mi.ras.ru, przhijal@mccme.ru`